



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books		
Search	PubMed	▼	for					Go	Clear	
		Limits	Preview/Index	History	Clipboard	Details				
		Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order

Entrez
PubMed

☐ 1: Mol Microbiol 1994 Sep;13(6):1093-100

[Related Articles, Books, LinkOut](#)

Anthrax toxin lethal factor contains a zinc metalloprotease consensus sequence which is required for lethal toxin activity.

Klimpel KR, Arora N, Leppla SH.

PubMed
Services

Laboratory of Microbial Ecology, National Institute of Dental Research, National Institutes of Health, Bethesda, Maryland 20892.

Related
Resources

Comparison of the anthrax toxin lethal factor (LF) amino acid sequence with sequences in the Swiss protein database revealed short regions of similarity with the consensus zinc-binding site, HEXXH, that is characteristic of metalloproteases. Several protease inhibitors, including bestatin and captopril, prevented intoxication of macrophages by lethal toxin. LF was fully inactivated by site-directed mutagenesis that substituted Ala for either of the residues (H-686 and H-690) implicated in zinc binding. Similarly, LF was inactivated by substitution of Cys for E-687, which is thought to be an essential part of the catalytic site. In contrast, replacement of E-720 and E-721 with Ala had no effect on LF activity. LF bound ⁶⁵Zn both in solution and on protein blots. The ⁶⁵Zn binding was reduced for several of the LF mutants. These data suggest that anthrax toxin LF is a zinc metalloprotease, the catalytic function of which is responsible for the lethal activity observed in cultured cells and in animals.

PMID: 7854123 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Sort	▼	Save	Text	Clip Add	Order
---------	----------	---	------	---	------	------	----------	-------

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)